

**Studies on Bacterial Terminal Quinol Oxidases from
JAPAN since
1928**

(June 17, 1999; 117 papers including 9 reviews, 9 Ph.D. theses and 10 studies (#) carried out abroad).

Comments should be sent to: T. Mogi (mogi@biol.s.u-tokyo.ac.jp).

Corresponding authors as relevant for each paper:

Yoshihiro Fukumori: fukumor@kenroku.ipc.kanazawa-u.ac.jp

Hisako Hirayama: hirayamah@jamstec.go.jp

Hiroshi Hori: hori@bpe.es.osaka-u.ac.jp

Masayasu Inoue: masainoue@msic.med.osaka-cu.ac.jp

Koreaki Ito: kito@virus.kyoto-u.ac.jp

HidekiKandori: kandori@photo2.biophys.kyoto-u.ac.jp

Chiaki Kato: katoc@jamstec.go.jp

Kiyoshi Kita: kitak@m.u-tokyo.ac.jp

Teizo Kitagawa: teizo@ims.ac.jp

Kazunobu Matsushita: kazunobu@agr.yamaguchi-u.ac.jp

Hideto Miyoshi: miyoshi@kais.kyoto-u.ac.jp

Tatsushi Mogi: mogi@biol.s.u-tokyo.ac.jp

Isao Morishima: morisima@mds.moleng.kyoto-u.ac.jp

Yutaka Orii: yorii@jade.dti.ne.jp [Retired, March 1998]

Junshi Sakamoto: sakamoto@bse.kyutech.ac.jp

Nobuhito Sone: sone@bse.kyutech.ac.jp

Tsuyoshi Sugio: sugio@cc.okayama-u.ac.jp

Motonari Tsubaki: tsubaki@sci.himeji-tech.ac.jp

Tsutomu Unemoto: unemoto@p.chiba-u.ac.jp

Isamu Yabe: yabe@bio.t.u-tokyo.ac.jp

Tateo Yamanaka: FAX, [81]-3-3293-7572

Note that the papers are in reverse chronological order!

Y. Yamazaki, H. Kandori, and T. Mogi (1999) Effects of subunit I mutations on redox-linked conformational changes of the Escherichia coli bo-type ubiquinol oxidase revealed by Fourier-transform infrared spectroscopy. *J. Biochem. (Tokyo)* 126, 194-199

Y. Yamazaki, H. Kandori, and T. Mogi (1999) Fourier-transform infrared studies on conformational changes of bd-type ubiquinol oxidase from Escherichia coli upon photoreduction of the redox metal centers. *J. Biochem. (Tokyo)* 125, 1131-1136

M. Tsubaki, T. Mogi, and H. Hori (1999) Azide- and cyanide-bindings to the Escherichia coli bd-type ubiquinol oxidase studied by visible absorption, EPR and FTIR spectroscopies. *J. Biochem. (Tokyo)* 126, in press

M. Tsubaki, T. Mogi, and H. Hori (1999) Fluoride-binding to the Escherichia coli bd-type ubiquinol oxidase studied by visible absorption and EPR spectroscopies. *J. Biochem. (Tokyo)* 126, 98-103

M. Tsubaki, T. Mogi, and H. Hori (1999) Fourier-transform infrared studies on azide binding to the binuclear center of the Escherichia coli bo-type ubiquinol oxidase. *FEBS Lett.* 449, 191-195

N. Sone, S. Tsukita and J. Sakamoto (1999) Direct correlation between proton translocation and growth yield: An analysis with a wild and mutant-type respiratory chains of Bacillus stearothermophilus. *J. Biosci. Bioeng.* 87, 495-499

J. Sakamoto, E. Koga, T. Mizuta, C. Sato, S. Noguchi, and N. Sone (1999) Gene structure and quinol oxidase activity of a cytochrome bd-type oxidase from Bacillus stearothermophilus. *Biochim. Biophys. Acta* 1411, 147-158

T. Mogi, M. Sato-Watanabe, H. Miyoshi, and Y. Orii (1999) Role of a bound ubiquinone on reactions of the Escherichia coli cytochrome bo with ubiquinol and dioxygen. *FEBS Lett.*, in press

H. Miyoshi, K. Takegami, K. Sakamoto, T. Mogi, and H. Iwamura (1999) Characterization of the ubiquinol oxidation sites in cytochromes bo and bd from Escherichia coli using aurachin C analogues. *J. Biochem. (Tokyo)* 125, 138-142

T. Kobayashi, and K. Ito (1999) Respiratory chain strongly oxidizes the

CXXC motif of DsbB in the Escherichia coli disulfide bond formation pathway. *EMBO J.* 18, 1192-1198

K. Kobayashi, S. Tagawa, and T. Mogi (1999) Pulse radiolysis studies on electron transfer processes in cytochrome bd-type ubiquinol oxidase from Escherichia coli. *Biochemistry* 38, 5913-5917

106)# L. Winstedt, K. I. Yoshida, Y. Fujita and C. von Wachenfeldt (1998) Cytochrome bd biosynthesis in Bacillus subtilis: Characterization of the cydABCD operon. *J. Bacteriol.* 180, 6571-6580

105) H. Tamegai, C. Kato, and H. Horikoshi (1998) Pressure-regulated respiratory system in barotolerant bacterium, Shewanella sp. strain DSS12. *J. Biochem. Mol. Biol. Biophys.* 1, 213-220

104) M. Sato-Watanabe, T. Mogi, H. Miyoshi, and Y. Anraku (1998c) Cooperation of two quinone binding sites in the oxidation of substrates by cytochrome bo. In: *Oxygen Homeostasis and Its Dynamics* (eds. Y. Ishimura, H. Shimada, M. Suematsu), pp.24-32. Springer-Verlag, Tokyo [Review]

103) M. Sato-Watanabe, T. Mogi, H. Miyoshi, and Y. Anraku (1998b) Isolation and characterizations of quinone analogue-resistant mutants of bo-type ubiquinol oxidase from Escherichia coli. *Biochemistry* 37, 12744-12752

102) M. Sato-Watanabe, T. Mogi, H. Miyoshi, and Y. Anraku (1998a) Characterization and functional role of the QH site of bo-type quinol oxidase from Escherichia coli. *Biochemistry* 37, 5356-5361

101) K. Sakamoto, H. Miyoshi, M. Ohshima, K. Kuwabara, K. Kano, T. Akagi, T. Mogi, and H. Iwamura (1998) Role of isoprenyl tail of ubiquinone in reaction with respiratory enzymes: Studies with bovine heart mitochondrial complex I and Escherichia coli bo-type ubiquinol oxidase. *Biochemistry* 37, 15106-15113

100) M. H. Qureshi, C. Kato and K. Horikoshi (1998) Purification of a ccb-type quinol oxidase specifically induced in a deep-sea barophilic bacterium, Shewanella sp. strain DB-172F. *Extremophiles* 2, 93-99

99) T. Mogi, M. Tsubaki, H. Hori, M. Miyoshi, H. Nakamura, and Y. Anraku (1998b) Two terminal quinol oxidase families in Escherichia coli: Variations on molecular machinery for dioxygen reduction. *J. Biochem. Mol. Biol. Biophys.* 2, 79-110 [Review]

98) T. Mogi, J. Minagawa, T. Hirano, M. Sato-Watanabe, M. Tsubaki, T.

- Uno, H. Hori, H. Nakamura, Y. Nishimura, and Y. Anraku (1998a) Substitutions of conserved aromatic amino acid residues in subunit I perturb the metal centers of the Escherichia coli bo-type ubiquinol oxidase. *Biochemistry* 37, 1632-1639
- 97) T. Kuroda, N. Okuda, N. Saitoh, Y. Terasaki, H. Anazawa, A. Hirata, T. Mogi, T. Hiyama, I. Kusaka, T. Tsuchiya, and I. Yabe (1998) Patch-clamp studies on ion pumps of the cytoplasmic membrane of Escherichia coli: Formation, preparation and utilization of giant vacuole-like structures consisting of everted cytoplasmic membranes. *J. Biol. Chem.* 273, 16897-16904
- 96) H. Hirayama, H. Takami, A. Inoue, and K. Horikoshi (1998) Isolation and characterization of toluene-sensitive mutants from *Pseudomonas putida* IH-2000. *FEMS Microbiol. Lett.* 169, 219-225
- 95) H. Yu, E. F. Sato, K. Nagata, M. Nishikawa, M. Kashiba, T. Arakawa, K. Kobayashi, T. Tamura and M. Inoue (1997) Oxygen-dependent regulation of the respiration and growth of Escherichia coli by nitric oxide. *FEBS Lett.* 409, 161-165
- 94) A. Yaginuma, S. Tsukita, J. Sakamoto and N. Sone (1997) Characterization of two terminal oxidases in *Bacillus brevis* and efficiency of energy conservation of the respiratory chain. *J. Biochem. (Tokyo)* 122, 969-976
- 93) A. Uchida, T. Kusano, T. Mogi, Y. Anraku, and N. Sone (1997) Expression of the Escherichia coli bo-type ubiquinol oxidase with a chimeric subunit II having the CuA-cytochrome c domain from the thermophilic *Bacillus caa3*-type cytochrome c oxidase. *J. Biochem. (Tokyo)* 122, 1004-1009
- 92) M. Tsubaki, H. Hori, and T. Mogi (1997b) Glutamate-286 mutants of cytochrome bo-type ubiquinol oxidase from Escherichia coli: Influence of mutations on the binuclear metal center structure revealed by FT-IR and EPR spectroscopies. *FEBS Lett.* 416, 247-250
- 91) M. Tsubaki, K. Matsushita, O. Adachi, S. Hirota, T. Kitagawa and H. Hori (1997a) Resonance Raman, Infrared, and EPR investigation on the binuclear site structure of the heme-copper ubiquinol oxidases from *Acetobacter acetii*: Effect of the heme peripheral formyl group substitution. *Biochemistry* 36, 13034-13042
- 90) M. Sato-Watanabe (1997) Studies on the ubiquinol oxidation site of the cytochrome bo-type ubiquinol oxidase in Escherichia coli. Ph. D. thesis, University of Tokyo [Thesis 9]

- 89) K. Saiki, T. Mogi, M. Tsubaki, H. Hori, and Y. Anraku (1997) Exploring subunit-subunit interactions in the *Escherichia coli* bo-type ubiquinol oxidase by extragenic suppressor mutation analysis. *J. Biol. Chem.* 272, 14721-14726
- 88) Y. Nogami, T. Maeda, A. Negishi and T. Sugio (1997) Inhibition of sulfur oxidizing activity by nickel iron in *Thiobacillus thiooxidans* NB1-3 isolated from the corroded concrete. *Biosci. Biotech. Biochem.* 61, 1373-1375
- 87) H. Nakamura, K. Saiki, T. Mogi, and Y. Anraku (1997) Assignment and functional roles of the *cyo*ABCDE gene products required for the *Escherichia coli* bo-type ubiquinol oxidase. *J. Biochem. (Tokyo)* 122, 415-421
- 86) T. Kobayashi, S. Kishigami, M. Sone, H. Inokuchi, T. Mogi, and K. Ito (1997) Respiratory chain is required to maintain oxidized states of the DsbA-DsbB disulfide bond formation system in aerobically growing *Escherichia coli* cells. *Proc. Natl. Acad. Sci. USA* 94, 11857-11862
- 85) M. Kawasaki, T. Mogi, and Y. Anraku (1997) Substitutions of charged amino acid residues conserved in subunit I perturb the redox metal centers of the *Escherichia coli* bo-type ubiquinol oxidase. *J. Biochem. (Tokyo)* 122, 422-429
- 84) C. Kato, L. Li, H. Tamegai, M. Smorawinska, and K. Horikoshi (1997b) A pressure-regulated gene cluster in deep-sea adapted bacteria with reference to its distribution. *Recent Res. Devel. Aggricul. Biol. Chem.* 1, 25-32
- 83) C. Kato, A. Ikegami, M. Smorawinska, R. Usami and K. Horikoshi (1997a) Structure of genes in a pressure-regulated operon and adjacent regions from a barotolerant bacterium strain DSS12. *J. Marine Biotech.* 5, 210-218
- 82) T. Hirano, T. Mogi, M. Tsubaki, H. Hori, Y. Orii and Y. Anraku (1997) A novel chloride-binding site modulates the heme-copper binuclear center in the *Escherichia coli* bo-type ubiquinol oxidase. *J. Biochem. (Tokyo)* 122, 430-437
- 81) M. Tsubaki, T. Mogi, H. Hori, M. Sato-Watanabe, and Y. Anraku (1996) Infrared and EPR studies on cyanide-binding to the heme-copper binuclear center of cytochrome bo-type ubiquinol oxidase from *Escherichia coli*: Release of a CuB-cyano complex in the partially reduced states. *J. Biol. Chem.* 271, 4107-4022

- 80) K. Sakamoto, H. Miyoshi, K. Takegami, T. Mogi, Y. Anraku, and H. Iwamura (1996) Probing substrate binding site of the *Escherichia coli* quinol oxidases using synthetic ubiquinol analogues based upon their electron-donating efficiency. *J. Biol. Chem.* 271, 29897-29902
- 79) J. Sakamoto, A. Matsumoto, K. Oobuchi and N. Sone (1996) Cytochrome bd-type quinol oxidase in a mutant of *Bacillus stearothermophilus* deficient in *caa3*-type cytochrome c oxidase. *FEMS Microbiol. Lett.* 143, 151-158
- 78) K. Saiki, T. Mogi, and Y. Anraku (1996b) Probing a role of subunit IV of the *Escherichia coli* bo-type ubiquinol oxidase by deletion and cross-linking analyses. *J. Biol. Chem.* 271, 15336-15340
- 77) K. Saiki (1996a) Discovery and reaction mechanism of the heme O synthase in *Escherichia coli*. Ph. D. thesis, University of Tokyo [Thesis 8]
- 76) C. Kato, H. Tamegai, A. Ikegami, R. Usami and K. Horikoshi (1996) Open reading frame 3 of the barotolerant bacterium strain DSS12 is complementary with *cydD* in *Escherichia coli*: *cydD* functions are required for cell stability at high pressure. *J. Biochem. (Tokyo)* 120, 301-305
- 75) H. Hori, M. Tsubaki, T. Mogi, and Y. Anraku (1996) EPR study of NO complex of bd-type ubiquinol oxidase from *Escherichia coli*. The proximal ligand of heme d is a nitrogenous amino acid residue. *J. Biol. Chem.* 271, 9254-9258
- 74) M. Tsubaki, H. Hori, T. Mogi, and Y. Anraku (1995) Cyanide-binding site of bd-type ubiquinol oxidase from *Escherichia coli*. *J. Biol. Chem.* 270, 28565-28569
- 73) M. Sato-Watanabe, S. Itoh, T. Mogi, K. Matsuura, H. Miyoshi, and Y. Anraku (1995) Stabilization of a semiquinone radical at the high affinity quinone binding site of the *Escherichia coli* bo-type ubiquinol oxidase. *FEBS Lett.* 374, 265-269
- 72) Y. Orii, T. Mogi, M. Sato-Watanabe, T. Hirano, and Y. Anraku (1995) Facilitated intramolecular electron transfer in the *Escherichia coli* bo-type ubiquinol oxidase requires chloride. *Biochemistry* 34, 1127-1132
- 71) T. Mogi, T. Hirano, H. Nakamura, Y. Anraku, and Y. Orii (1995) CuB promotes both binding and reduction of dioxygen at the heme-copper binuclear center in the *Escherichia coli* bo-type ubiquinol oxidase. *FEBS Lett.* 370, 259-263

- 70) S. Hirota, T. Mogi, T. Ogura, Y. Anraku, R. B. Gennis, and T. Kitagawa (1995) Resonance Raman study on axial ligands of heme irons in cytochrome bd-type ubiquinol oxidase from *Escherichia coli*. *Biospectroscopy* 1, 305-311
- 69) K. Denda, T. Mogi, Y. Anraku, T. Yamanaka, and Y. Fukumori (1995) Characterization of chimeric heme-copper respiratory oxidases using subunits I of *Escherichia coli* cytochrome bo and *Halobacterium salinarium* cytochrome aa₃. *Biochem. Biophys. Res. Commun.* 217, 428-436
- 68) T. Uno, T. Mogi, M. Tsubaki, Y. Nishimura, and Y. Anraku (1994) Resonance Raman and Fourier-transform infrared studies on the subunit I histidine mutants of the cytochrome bo complex in *Escherichia coli*: Molecular structure of redox centers. *J. Biol. Chem.* 269, 11912-11920
- 67) M. Tsubaki, T. Mogi, H. Hori, T. Ogura, S. Hirota, T. Kitagawa, and Y. Anraku (1994) Molecular structure of redox metal centers of the cytochrome bo complex from *Escherichia coli*: Spectroscopic characterizations of the subunit I histidine mutant oxidases. *J. Biol. Chem.* 269, 30861-30868
- 66) M. Sato-Watanabe, T. Mogi, T. Ogura, T. Kitagawa, H. Miyoshi, H. Iwamura, and Y. Anraku (1994b) Identification of a novel quinone binding site in the cytochrome bo complex from *Escherichia coli*. *J. Biol. Chem.* 269, 20908-20912
- 65) M. Sato-Watanabe, T. Mogi, H. Miyoshi, H. Iwamura, K. Matsushita, O. Adachi, and Y. Anraku (1994a) Structure-function studies on the ubiquinol oxidation site of the cytochrome bo complex from *Escherichia coli* using p-benzoquinones and substituted phenols. *J. Biol. Chem.* 269, 28899-28907
- 64) K. Saiki, T. Mogi, M. Ishizuka, and Y. Anraku (1994) An *Escherichia coli* cyoE gene homologue in thermophilic *Bacillus* PS3 encodes a thermotolerant heme O synthase. *FEBS Lett.* 351, 385-388
- 63) S. Ozawa, Y. Watanabe and I. Morishima (1994d) Spectroscopic characterization of peroxo-iron(III) complexes. The first model for a reaction intermediate of cytochrome d. *Inorg. Chem.* 33, 306-313
- 62) S. Ozawa, Y. Watanabe and I. Morishima (1994c) NMR studies of p-actin radical complexes of iron(III) and oxoiron(IV) chlorins. Models for reaction intermediates of chlorin-containing heme enzymes. *J. Am. Chem. Soc.* 116, 5832-5838

- 61) S. Ozawa, Y. Watanabe, S. Nakashima, T. Kitagawa and I. Morishima (1994b) Preparation and characterization of oxoiron(IV) chlorin complexes as the first models for a reaction intermediate in the catalytic cycle of cytochrome d. *J. Am. Chem. Soc.* 116, 634-641
- 60) S. Ozawa (1994a) The electronic structures and functions of the reaction intermediates in hdroporphyrin-containing heme enzymes. Ph. D. thesis, Kyoto University [Thesis 7]
- 59) Y. Orii, T. Mogi, M. Kawasaki, and Y. Anraku (1994) Facilitated intramolecular electron transfer in the cytochrome bo-type ubiquinol oxidase initiated upon reaction of the fully-reduced enzyme with dioxygen. *FEBS Lett.* 352, 151-154
- 58) T. Mogi, H. Nakamura, and Y. Anraku (1994b) Molecular structure of a heme-copper redox center of the *Escherichia coli* ubiquinol oxidase: Evidence and model. *J. Biochem. (Tokyo)* 116, 741-747 [Review]
- 57) T. Mogi, K. Saiki, and Y. Anraku (1994a) Biosynthesis and functional role of haem O and haem A. *Mol. Microbiol.* 14, 391-398 [Review]
- 56) S. Hirota, T. Mogi, T. Ogura, T. Hirano, Y. Anraku, and T. Kitagawa (1994b) Observation of the Fe-O₂ and FeIV=O Raman bands in dioxygen reduction by cytochrome bo-type ubiquinol oxidase from *Escherichia coli*. *FEBS Lett.* 352, 67-70
- 55) S. Hirota (1994a) Resonance Raman studies of Fe-ligand vibrations of oxygen binding heme proteins and structure-function relationship of terminal oxidases. Ph. D. thesis, Graduate University for Advanced Studies [Thesis 6]
- 54) M. Tsubaki, T. Uno, H. Hori, T. Mogi, Y. Nishimura, and Y. Anraku (1993b) Heme D axial ligand of the cytochrome bd complex from *Escherichia coli*. *FEBS Lett.* 335, 13-17
- 53) M. Tsubaki, T. Mogi, Y. Anraku, and H. Hori (1993a) Structure of heme-copper binuclear center of the cytochrome bo complex of *Escherichia coli*: EPR and Fourier-transform infrared spectroscopic studies. *Biochemistry* 32, 6065-6072
- 52) K. Saiki, T. Mogi, H. Hori, M. Tsubaki, and Y. Anraku (1993b) Identification of the functional domains in heme O synthase: Site-directed mutagenesis studies on the *cyoE* gene of the cytochrome bo operon in *Escherichia coli*. *J. Biol. Chem.* 268, 26927-26934

- 51) K. Saiki, T. Mogi, K. Ogura, and Y. Anraku (1993a) In vitro heme O synthesis by the *cyoE* gene product from *Escherichia coli*. *J. Biol. Chem.* 268, 26041-26045
- 50) T. Miyoshi-Akiyama, M. Hayashi and T. Unemoto (1993) Purification and properties of cytochrome *bo*-type ubiquinol oxidase from a marine bacterium *Vivrio alginolyticus*. *Biochim. Biophys. Acta* 1141, 283-287
- 49) M. Fukaya, K. Tayama, T. Tamaki, H. Ebisuya, H. Okumura, Y. Kawamura, S. Horinouchi, and T. Beppu (1993) Characterization of a cytochrome *a1* that functions as a ubiquinol oxidase in *Acetobacter aceti*. *J. Bacteriol.* 175, 4307-4314
- 48) K. Saiki, T. Mogi, and Y. Anraku (1992) Heme O biosynthesis in *Escherichia coli*: The *cyoE* gene in the cytochrome *bo* operon encodes a protoheme IX farnesyltransferase. *Biochem. Biophys. Res. Commun.* 189, 1491-1497
- 47) S. Ozawa, Y. Watanabe and I. Morishima (1992) Preparation and characterization of a novel oxoiron(IV) chlorin pi-cation radical complex. The first model for compound I of chlorin-containing heme enzymes. *Inorg. Chem.* 31, 4042-4043
- 46) J. Minagawa, T. Mogi, R. B. Gennis, and Y. Anraku (1992b) Identification of heme and copper ligands in subunit I of the cytochrome *bo* complex in *Escherichia coli*. *J. Biol. Chem.* 267, 2096-2104
- 45) J. Minagawa (1992a) Molecular structure of the redox reaction center of the cytochrome *bo* complex in *Escherichia coli*. Ph. D. thesis, University of Tokyo [Thesis 5]
- 44) K. Matsushita, H. Ebisuya, M. Ameyama and O. Adachi (1992b) Change of the terminal oxidase from cytochrome *a1* in shaking cultures to cytochrome *o* in static cultures of *Acetobacter aceti*. *J. Bacteriol.* 174, 122-129
- 43) K. Matsushita, H. Ebisuya and O. Adachi (1992a) Homology in the structure and the prosthetic groups between two different terminal ubiquinol oxidases, cytochrome *a1* and cytochrome *o*, of *Acetobacter aceti*. *J. Biol. Chem.* 267, 24748-24753
- 42) H. Nakamura (1991) Studies on the *Escherichia coli* Cytochrome *b562-o* Terminal Oxidase Complex in the Aerobic Respiratory Chain. Ph. D. thesis, University of Tokyo [Thesis 4]
- 41) H. Nakamura, I. Yamato, Y. Anraku, L. Lemieux, and R. B. Gennis

(1990) Expression of *cyoA* and *cyoB* demonstrates that the CO-binding heme component of the *Escherichia coli* cytochrome *o* complex is in subunit I. *J. Biol. Chem.* 265, 11193-11197

40) J. Minagawa, H. Nakamura, I. Yamato, T. Mogi, and Y. Anraku (1990) Transcriptional regulation of the cytochrome *b562-o* complex in *Escherichia coli*. Gene expression and molecular characterization of the promoter. *J. Biol. Chem.* 265, 11198-11203

39) K. Matsushita, E. Shinagawa, O. Adachi and M. Ameyama (1990) Cytochrome *a1* of *Acetobacter aceti* is a cytochrome *ba* functioning as ubiquinol oxidase. *Proc. Natl. Acad. Sci. USA* 87, 9863-9867

38)# R. M. Lorence, K. Carter, R. B. Gennis, K. Matsushita and H. R. Kaback (1988) Trypsin proteolysis of the cytochrome *d* complex of *Escherichia coli* selectively inhibits ubiquinol oxidase activity while not affecting N,N,N',N'-tetramethyl-p-phenylenediamine oxidase activity. *J. Biol. Chem.* 263, 5271-5276

37) Y. Anraku (1988) Bacterial electron transport chains. *Annu. Rev. Biochem.* 57, 101-132 [Review]

36) K. Matsushita, E. Shinagawa, O. Adachi and M. Ameyama (1987) Purification, characterization and reconstitution of cytochrome *o*-type oxidase from *Gluconobacter suboxydans*. *Biochim. Biophys. Acta* 894, 304-312

35) A. Hata-Tanaka, K. Matsuura, S. Itoh, and Y. Anraku (1987) Electron flow and heme-heme interaction between cytochrome *b-558*, *b595* and *d* in a terminal oxidase of *Escherichia coli*. *Biochim. Biophys. Acta* 893, 289-295

34)# M. J. Costello, J. Escaig, K. Matsushita, P. V. Viitanen, D. R. Menick and H. R. Kaback (1987) Purified lac permease and cytochrome *o* oxidase are functional as monomers. *J. Biol. Chem.* 262, 17072-17082

33) Y. Anraku (1987) Biochemistry and molecular biology of the *Escherichia coli* aerobic respiratory chain. In: *Cytochrome Systems* (Eds., S. Papa, B. Chance, and L. Ernster), Plenum Publ.

32) Y. Anraku, and R. B. Gennis (1987) The aerobic respiratory chain of *Escherichia coli*. *Trends Biochem. Sci.* 12, 262-266 [Review]

31)# C. C. Moser, K. Matsushita, D. E. Robertson, H. R. Kaback and P. L. Dutton (1986) Kinetics of electron transfer in reaction center-

cytochrome o proteoliposomes. FEBS Lett. 194, 115-120

30)# K. Matsushita and H. R. Kaback (1986b) D-Lactate oxidation and generation of the proton electrochemical gradient in membrane vesicles from *Escherichia coli* GR19N and in proteoliposomes reconstituted with purified D-lactate dehydrogenase and cytochrome o oxidase. *Biochemistry* 25, 2321-2327

29)# K. Matsushita, L. Patel and H. R. Kaback (1986a) Purification and reconstitution of the cytochrome o-type oxidase from *Escherichia coli*. *Methods. Enzymol.* 126, 113-122 [Review]

28) K. Konishi, M. Ouchi, K. Kita and I. Horinouchi (1986) Purification and properties of a cytochrome b560-d complex, a terminal oxidase of the aerobic respiratory chain of *Photobacterium phosphoreum*. *J. Biochem. (Tokyo)* 99, 1227-1236

27) K. Kita, K. Konishi and Y. Anraku (1986) Purification and properties of two terminal oxidase complexes of *Escherichia coli* aerobic respiratory chain. *Methods. Enzymol.* 126, 94-113 [Review]

26)# T. Hamamoto and M. Montal (1986) Functional reconstitution of bacterial cytochrome oxidases in planar lipid bilayers. *Methods. Enzymol.* 126, 123-138 [Review]

25) T. Uno, Y. Nishimura, M. Tsuboi, K. Kita, and Y. Anraku (1985) Resonance Raman study of cytochrome b562-o complex, a terminal oxidase of *Escherichia coli* in its ferric, ferrous, and CO-ligated states. *J. Biol. Chem.* 260, 6755-6760

24) K. Kita, H. Murakami, H. Ohya and Y. Anraku. (1985) Quantitative determination of cytochromes in the aerobic respiratory chain of *Escherichia coli* by high-performance liquid chromatography and its application to analysis of mitochondrial cytochromes. *Biochem. Int.* 10, 319-326

23) A. Hata, Y. Kirino, K. Matsuura, S. Itoh, T. Hiyama, K. Konishi, K. Kita, and Y. Anraku (1985b) Assignment of ESR signals of *Escherichia coli* terminal oxidase complexes. *Biochim. Biophys. Acta* 810, 62-72

22) A. Hata (1985a) Electronic states of terminal oxidase cytochrome b558-d complex in *Escherichia coli*. Ph. D. thesis, University of Tokyo [Thesis 3]

21)# T. Hamamoto, N. Carrasco, K. Matsushita, H. R. Kaback and M. Montal (1985) Direct measurement of the electrogenic activity of o-type

cytochrome oxidase from *Escherichia coli* reconstituted into planar lipid bilayers. *Proc. Natl. Acad. Sci. USA* 82, 2570-2573

20)# K. Matsushita, L. Patel, and H. R. Kaback (1984) Cytochrome o type oxidase from *Escherichia coli*. Characterization of the enzyme and mechanism of electrochemical gradient generation. *Biochemistry* 23, 4703-4714

19) K. Kita, K. Konishi, and Y. Anraku (1984b) Terminal oxidases of *Escherichia coli* aerobic respiratory chain. II. Purification and properties of cytochrome b558-d complex from cells grown with limited oxygen and evidence of branched electron-carrying systems. *J. Biol. Chem.* 259, 3375-3381

18) K. Kita, K. Konishi, and Y. Anraku (1984a) Terminal oxidases of *Escherichia coli* aerobic respiratory chain. I. Purification and properties of cytochrome b562-o complex from cells in the early exponential phase of aerobic growth. *J. Biol. Chem.* 259, 3368-3374

17)# K. Matsushita, L. Patel, R. B. Gennis, and H. R. Kaback (1983) Reconstitution of active transport in proteoliposomes containing cytochrome o oxidase and lac carrier protein purified from *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 80, 4889-4893

16) K. Konishi (1983) Purification and properties of cytochrome b558-d in the respiratory chain of *Escherichia coli*. Ph. D. thesis, University of Tokyo (in Japanese) [Thesis 2]

15) K. Kita, M. Kasahara, and Y. Anraku. (1982) Formation of a membrane potential by reconstituted liposomes made with cytochrome b562-o complex, a terminal oxidase of *Escherichia coli* K12. *J. Biol. Chem.* 257, 7933-7935

14) K. Kita, and Y. Anraku. (1981) Composition and sequence of b cytochromes in the respiratory chain of aerobically grown *Escherichia coli* K-12 in the early exponential phase. *Biochem. Int.* 2, 105-112

13) K. Kita (1980) Studies on cytochrome components in the respiratory chain of *Escherichia coli*. Ph. D. thesis, University of Tokyo (in Japanese) [Thesis 1]

12) H. Watanabe, Y. Kamita, T. Nakamura, A. Takimoto, and T. Yamanaka (1979) The terminal oxidase of *Photobacterium phosphoreum*. A novel cytochrome. *Biochim. Biophys. Acta* 547, 70-78

11) A. Ishida and N. Shiraishi (1978) Development of hemoproteins and

respiratory activities of *Escherichia coli* grown at high and low aeration. *J. Gen. Appl. Microbiol.* 24, 139-141

10) Y. Inoue (1976) Presence of cytochrome d in respiratory particles of *Streptomyces griseus* and its function in the terminal oxidation system. *Bot. Mag. Tokyo* 89, 183-190

9) Y. Inoue (1973) The metabolism of *Streptomyces griseus*. XI. Further observation on the electron transfer system of the *Streptomyces griseus* respiratory particle. *Bot. Mag. Tokyo* 86, 121-132

8) A. Ishida and S. Hino (1972) Effect of oxygen on cytochrome pattern and heme synthesis in *Escherichia coli*. *J. Gen. Appl. Microbiol.* 18, 25-237

7) S. Hino and M. Maeda (1966) Effect of oxygen on the development of respiratory activity in *Escherichia coli*. *J. Gen. Appl. Microbiol.* 12, 247-265

6) T. Oka and K. Arima (1965) Cyanide resistance in *Achromobacter*. II. Mechanism of cyanide resistance. *J. Bacteriol.* 90, 744-747

5) K. Arima and T. Oka (1965) Cyanide resistance in *Achromobacter*. I. Induced formation of cytochrome a₂, and its role in cyanide-resistant respiration. *J. Bacteriol.* 90, 734-743

4) S. Yamaguchi (1937) Einige Untersuchungen über das Cytochrom der Bakterien. *Bot. Mag. Tokyo* 51, 457-461

3) A. Fujita and T. Kodama (1934) Untersuchungen über Atmung und Gärung pathogener Bakterien. III. Mitteilung: Über Cytochrom und das sauerstoffübertragende Ferment, sowie die Atmungshemmung der pathogenen Bakterien durch CO und HCN. *Biochim. Z.* 273, 186-197

2) H. Tamiya and S. Yamaguchi (1933) Systematische Untersuchungen über das Cytochromspektrum von verschiedenen Mikroorganismen. *Acta Phytochim.* 7, 233-244

1) H. Yaoi and H. Tamiya (1928) On the respiratory pigment, cytochrome, in bacteria. *Proc. Imp. Acad. Japan* 4, 436-439

Related Enzymes (8 papers as of June 14, 1999).

S. Tsukita, S. Koyanagi, K. Nagata, H. Koizuka, H. Akashi, T. Shimoyama, T. Tamura, and N. Sone (1999) Characterization of a cb-type cytochrome c oxidase from *Helicobacter pylori*. *J. Biochem. (Tokyo)* 125, 194-201

J. Sakamoto, A. Hayakawa, T. Uehara, S. Noguchi, and N. Sone (1999) Cloning of *Bacillus stearothermophilus* ctaA and heme A synthesis with the CtaA protein produced in *Escherichia coli*. *Biosci. Biotechnol. Biochem.* 63, 96-103

T. Tanaka, M. Inoue, J. Sakamoto, and N. Sone (1996) Intra- and Inter-complex cross-linking of subunits in the quinol oxidase super-complex from thermophilic *Bacillus PS3*. *J. Biochem.* 119, 482-486

M. Yamada, K. Sumi, K. Matsushita, O. Adachi, and Y. Yamada (1993) Topological analysis of quinoprotein glucose dehydrogenase in *Escherichia coli* and its ubiquinone-binding site. *J. Biol. Chem.* 268, 12812-12817

K. Matsushita, Y. Takaki, E. Shinagawa, M. Ameyama, and O. Adachi (1992b) Ethanol oxidase respiratory chain of acetic acid bacteria. Reactivity with ubiquinone of pyrroloquinoline quinone-dependent alcohol dehydrogenases purified from *Acetobacter aceti* and *Gluconobacter suboxydans*. *Biosci. Biotech. Biochem.* 56, 304-310

K. Matsushita, K. Takahashi, M. Takahashi, M. Ameyama, and O. Adachi (1992a) Methanol and ethanol oxidase respiratory chains of the methylotrophic acetic acid bacterium, *Acetobacter methanolicus*. *J. Biochem. (Tokyo)* 111, 739-747

K. Matsushita, M. Nonobe, E. Shinagawa, O. Adachi, and M. Ameyama (1987) Reconstitution of pyrroloquinoline quinone-dependent D-glucose oxidase respiratory chain of *Escherichia coli* with cytochrome o oxidase; *J. Bacteriol.* 169, 205-209

T. Yamanaka and Y. Kamita (1981) Purification and some properties of a c-type cytochrome derived from *Photobacterium phosphoreum*. *J. Gen. Appl. Microbiol.* 27, 273-275